



Hackensack
Meridian Health

**WOMEN'S CENTER
SAVI SCOUT PRE-PROCEDURE
INSTRUCTIONS**



OP5212

70308-014CX (11-19-18) PAGE 1 OF 2

PATIENT LABEL

Savi Scout Reflector Placement: Left Right

Date of Procedure: _____ Arrival Time: _____ Appt Time: _____

1. All patients are responsible to get precertification and/or preauthorization for biopsies. Pre-registration will contact you at least 2 business days before your biopsy. If they do not reach you, please call the pre-registration desk at 732-897-7979.
2. Please take all prescribed medications. DO NOT stop your medication for Savi Scout Reflector Placement Procedure.
 - a. Are you allergic to Nickel? Yes No Don't Know
 - b. Do you have a defibrillator or cardiac pacemaker? Yes No
 - c. Are you older than 18 years old? Yes No
3. You may take Tylenol or Tylenol Extra Strength for pain relief.
4. You should eat a light meal the day of your biopsy.
5. You may drive yourself; however, if you need to take an anti-anxiety medication, you must have a driver to and from the hospital. We do not prescribe or dispense these medications,
6. Discharge Instructions (more detail will be provided at discharge):
7. If you have any questions contact the nurse in the Women's Center where you will be having your procedure:
Riverview Women's Center - (732) 530-2308 Bayshore Women's Center - (732) 497-1870

Additional Instructions: _____

CONFIRMED HEIGHT/WEIGHT PER EQUIPMENT REQUIREMENTS

Biopsy Instruction Packet given to Pt: Yes No MAILED/FAXED: _____ Date: _____

Instructions provided by: _____ Date: _____ Time: _____ AM / PM

Patient Signature: _____ Date: _____ Time: _____ AM / PM



Tylenol or Tylenol Extra Strength are acceptable pain relievers before and after procedure.

If you are on medications such as anti-coagulants (e.g., Coumadin) or anti-depressants (e.g., nardil) you must inform the nurse in Women's Center.

Some common products, including medications that contain aspirin or ibuprofen, which may impair blood clotting:

Advil	Arthritis Pain Formula	Darvon Compound	Fiorinal
Alka Seltzer	Ascriptin	Ecotrin	Midol
Aleve	Aspergum	Empirin	Nuprin
Anacin	Bayer- Time release	Excedrin	Percodan
Bufferin	Excedrin PM	Sine-Aid	

Nutritional supplements which may impair blood clotting:

Fish Oil, Garlic, Ginko-Biloba, Gotu Kola, Horse Chestnut, Vitamin C, Vitamin E and St. John's Wort